

In re Patent Application of:

OLSSON ET AL.

Serial No. 09/147,230

Filed: 2/9/99

A1
Concl

where N is the number of active carriers and $(X_{n,k})(Y_{n,k})$ is the unwrapped argument function for the nth carrier in the kth frame. --

[

Replace the paragraph beginning at page 6, line 4 through page 6, line 9, with the following rewritten paragraph:

-- Said slope of said argument function, α_k , may be estimated from the equation

$$\alpha_k = \frac{2}{n_2 - n_0} \left[\sum_{n=n_1+1}^{n_2} L(X_{n,k}) / (Y_{n,k}) - \sum_{n=n_0}^{n_1} L(X_{n,k})(Y_{n,k}) \right]$$

A2

where N is the number of active carriers, $(X_{n,k})(Y_{n,k})$ is the unwrapped argument function for the nth active carrier in the kth frame, indices n_0 and n_2 are the lower and upper limits respectively of the band and index n_1 divides the band into two equal parts. --

[

Replace the paragraph beginning at page 8, line 3 through page 8, line 7, with the following rewritten paragraph:

Said slope of said argument function, α_k , may be estimated from the equation

A3
Cott

$$\alpha_k = \frac{1}{N} \sum_n L \frac{(X_{n,k})(Y_{n,k})}{n}$$

In re Patent Application of:

OLSSON ET AL.

Serial No. 09/147,230

Filed: 2/9/99

A3
Coul

where N is the number of active carriers and $(X_{n,k})(Y_{n,k})$ is the unwrapped argument function for the nth carrier in the kth frame. --

[Replace the paragraph beginning at page 8, line 8 through page 8, line 13, with the following rewritten paragraph:

-- Said slope of said argument function, α_k , may be estimated from the equation

A4

$$\alpha_k = \frac{2}{n_2 - n_0} \left[\sum_{n=n_1+1}^{n_2} L(X_{n,k}) / (Y_{n,k}) - \sum_{n=n_0}^{n_1} L(X_{n,k})(Y_{n,k}) \right]$$

where N is the number of active carriers, $(X_{n,k})(Y_{n,k})$ is the unwrapped argument function for the nth active carrier in the kth frame, indices n_0 and n_2 are the lower and upper limits respectively of the band and index n_1 divides the band into two equal parts. --

[Paragraph beginning at page 11, line 19 through page 12, line 4, has been amended as follows:

-- The average slope, α_k , of the linear part of the argument function can be calculated, as shown in equation (1), or by some other standard method, using the unwrapped argument function of X_k for the kth frame

A5

$$\alpha_k = \frac{1}{N} \sum_n L \frac{(X_{n,k})(Y_{n,k})}{n} \dots\dots\dots(1)$$